

SS Ari: a shallow-contact close binary system

Liu L.^{1,2,3}, Qian S.-B.^{1,2,3}, He J.-J.^{1,2,3}, Zhang J.^{1,2,3} and Li L.-J.^{1,2,3}

ABSTRACT

Two CCD epochs of light minimum and a complete R light curve of SS Ari are presented. The light curve obtained in 2007 was analyzed with the 2003 version of the W-D code. It is shown that SS Ari is a shallow contact binary system with a mass ratio $q = 3.25$ and a degree of contact factor $f = 9.4\%(\pm 0.8\%)$. A period investigation based on all available data shows that there may exist two distinct solutions about the assumed third body. One, assuming eccentric orbit of the third body and constant orbital period of the eclipsing pair results in a massive third body with $M_3 = 1.73M_\odot$ and $P_3 = 87.0\text{yr}$. On the contrary, assuming continuous period changes of the eclipsing pair the orbital period of tertiary is 37.75yr and its mass is about $0.278M_\odot$. Both of the cases suggest the presence of an unseen third component in the system.

Subject headings: Stars: binaries : close – Stars: binaries : eclipsing – Stars: individuals (SS Ari) – Stars: evolution

1. Introduction

SS Ari, which was discovered as a variable star by Hoffmeister (1934) and then was recognized as a W Ursae Majoris type eclipsing binary system, has been paid more and more attention to due to its erratic period variations (Demircan & Selam 1993) since 1993. Many high accurate CCD times of minimum of it has been obtained in recent years. Five years have past since the last researches have been done by Kim et al. (2003). It is high time that we did some new studies to compare with the results found out by our predecessors.

¹National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences, P.O. Box 110, 650011 Kunming, P.R. China (e-mail: creator_ll.student@sina.com; LiuL@ynao.ac.cn)

²United Laboratory of Optical Astronomy, Chinese Academy of Sciences (ULOAC), 100012 Beijing, P. R. China

³Graduate School of the CAS, Beijing, P.R. China

Many researchers have presented its photoelectric light, such as Zhukov (1975), Kaluzny & Pojmanski (1984a, b), Rainger et al. (1992), Liu et al. (1993) and Kim et al. (2003). At the same time, spectroscopic observations for this system have been reported by Lu (1991), Rainger et al. (1992) and Kim et al. (2003). Furthermore, the period variabilities were studied by Braune (1970), Kaluzny & Pojmanski (1984a, b), Kurpinska-Winiarska & Zakrzewski (1990), Rainger et al. (1992), Demircan & Selam (1993), Kim et al. (1997), Kim et al. (2003). And so on.

Up to now, it is agreed that SS Ari is a sun-like W-type W UMa system with a spectral type in the interval F8V-G2V, mass parameters $M_1 = 0.4M_\odot$, $M_2 = 1.3M_\odot$. As Kim et al. (2003) pointed out, there are still many discordant matters among the previous investigations: diverse interpretations of the period activity, apparent intrinsic variability of the light curve, and disparate radial velocity curves. New times of minimum can validate the previous results about its period changes; new light curves can hope to make a contribution to the first two confusions. Those are the motivation of our current work.

2. New observations

SS Ari was observed on two nights (December 13, 14, 2007) with the PI512 TKB CCD photometric system attached to the 60cm reflecting telescope at the Yunnan Observatory in China. The R color system used is close to the standard UBVRI system. The effective field of view of the photometric system is 7×7 arcmin at the Cassegrain focus. The integration time for each image is 15s. The comparison star is BD+23°277. PHOT (measure magnitudes for a list of stars) of the aperture photometry package of IRAF was used to reduce the observed images. Through the observation we obtained a complete R light. By calculating the phase of the observations with Equation 2, the light curves are plotted (Figure 1) and the original data in the R band are listed in Table 1. In this figure, it is shown that the data are high quality and the light variation is typical of EW type. Since the lights around the minimum are symmetric, a parabolic fitting was used to determine the times of minimum light by the least square method. In all, our two new epochs of light minimum were obtained and listed in the last line of Table 2.

3. Orbital period variations

The orbital period of the W-type contact binary, SS Ari, has been studied by several authors. Braune (1970) first noted that the period of SS Ari is variable. And this conclusion

was confirmed later by Kaluzny & Pojmanski (1984b). They deemed that the period can be described by two possibilities, (i) continuous sine-like variation, (ii) change formed by several jumps. Kurpinska-Winiarska & Zakrzewski (1990) questioned the sine-like shape of the O-C curve of SS Ari, since the sine-like variation mainly results from the positions of the visual and photographic timings published before 1965. They forced to present the O-C observations by a parabolic fitting, claiming that the sine-like fitting was also acceptable if the O-C values of the timings published before 1965 were shifted by half a period. Later, the period changes of SS Ari were studied in detail by Demircan & Selam (1993); their results showed that the O-C diagram of the system varies as a cyclic oscillation of which the amplitude is $0.^d0398$ and the period is 44.8 yr. This phenomena was explained as light-time effect, which is caused by a hypothetical third body with $M_3 \sim 1.0M_\odot$. Since not any sign of the third body was seen in the spectrum of this binary, Lu(1991) pointed out that the hypothetical third body is too faint, most probably a binary just as in the case of XY Leo (Barden 1987), or a white dwarf. The latest researches were given by Kim et al. (2003). They postulated three possible reasons for the period variation. First, it was caused by cyclical magnetic activity or by a light-time effect due to a gravitationally bound third star. Second, it may be a secular period change because of mass transfer. Third, the real variations were more complicated. They concluded that there exist not only a third body ($m_3 = 1.96M_\odot, P_3 = 39.7\text{yr}$), but also a fourth body ($m_4 = 2.38M_\odot, P_4 = 88.2\text{yr}$).

3.1. Model I: A constant orbit superimposed a eccentric orbit fitting

Demircan & Selam gave their liner ephemeris:

$$MinI = 2444469.4790 + 0^d.40599144 \times E \quad (1)$$

After that investigation, some photoelectric and CCD times of light minimum have been published. All its available times of light minimum have been collected to look for the real period changes, including visual, photographic, photoelectric and CCD observations. Those were completely listed in Table 2. Based on all collected eclipse times, setting visual and photographic data as weight 1, meanwhile setting photoelectric and CCD data as weight 8, a new linear ephemeris was obtained:

$$MinI = 2444469.4879(\pm 0.0030) + 0^d.40598629(\pm 0.00000026) \times E. \quad (2)$$

The $(O - C)$ values with respect to the linear ephemeris are listed in the fifth column of Table 2. The corresponding $(O - C)$ diagram is displayed in Figure 2. It is no doubt that the general $(O - C)$ trend of SS Ari, shown in Figure 2, is a cyclic variation. Consider the

relationship of the two periods derived by Kim et al. (2003), an eccentric orbit and constant period fitting was used. The result is:

$$\begin{aligned} MinI &= 2444469.4169 + 0.40598629 \times E \\ &+ 0.0859 \cos 0^\circ.0046E + 0.0595 \sin 0^\circ.0046E \\ &+ 0.0023 \cos 0^\circ.0092E - 0.0205 \sin 0^\circ.0092E. \end{aligned} \quad (3)$$

The corresponding O-C curve is shown in Figure 2, where solid cycles refer to the photoelectric or CCD primary minima and open ones to the photoelectric or CCD secondary minima. Solid triangles denote visual or photographic primary minima, open triangles denote visual or photographic secondary minima; meanwhile, solid line represents an eccentric ephemeris variation. The corresponding residuals were drawn in Figure 3. From it we can see all photoelectric, CCD and most of visual, photographic timings' residuals are in a horizontal line, except some scatters. It accounts for that the line fits all photoelectric and CCD data very well, although the visual and photographic data show large scatters in the E interval 0-10000. By aid of relations (Kopal 1978),

$$\omega = 360^\circ P_e / T, \quad (4)$$

$$e' = 2 \sqrt{\frac{a_2^2 + b_2^2}{a_1^2 + b_1^2}}, \quad (5)$$

$$w' = \arctan \frac{(b_1^2 - a_1^2)b_2 + 2a_1b_1a_2}{(a_1^2 - b_1^2)a_2 + 2a_1b_1b_2}, \quad (6)$$

where P_e is the ephemeris period ($0.^d40598629$) and a_1, a_2, b_1, b_2 are the corresponding fitting coefficients ($a_1 = 0.0859, b_1 = 0.0595, a_2 = 0.0023, b_2 = -0.0205$), the period of the orbital oscillation was determined to be $T=87.0\text{yr}$ with an eccentricity $e' = 0.3948$. These may suggest that there exists a small-amplitude oscillation in the period changes, which can be explained by the presence of an unseen third body in the system. This unseen third body was first proposed by Demircan & Selam (1993) and later confirmed by Kim et al. (2003).

3.2. Model II: A quadratic change superimposed a cycle change fitting

However, since very small amount of the early data (before 1960), the photographic and visual times of light minimum showed a large scatter (up to 0.06 days). Those data not only do not contribute to form the general O-C trend of the binary system but also tend to mislead regarding the real period changes. Considered this, the key visual data in early years may be fallible, the simulation by eccentric orbit and constant period model may be incredible.

The same situation was met during the period analysis of AM Leo (Qian et al. 2005) and HL Aur (Qian et al. 2006), AP Leo (Qian et al. 2007). This situation suggests that some of the period variations pointed out by former investigators are hardly reliable. Therefore, only using those photoelectric and CCD times of minimum light for the present period study of SS Ari is necessary. The new linear ephemeris purely derived by photographic and CCD data is:

$$MinI = 2444469.4789(\pm 0.0071) + 0^d.40599166(\pm 0.00000053) \times E. \quad (7)$$

Based on these high accurate data, we give a quadratic fitting:

$$\begin{aligned} MinI = & 2444469.4988(\pm 0.0004) + 0.40598838(\pm 0.00000002) \times E \\ & - 1.72(\pm 0.01) \times 10^{-10} E^2. \end{aligned} \quad (8)$$

That fitting is shown in Figure 4, the dash line. Actually, only a quadratic simulation can not fit the data well. Having bethought of the doubtless cyclic oscillation in O-C diagram, we used a circular orbit and continuous period change form to fit these data, the result is:

$$\begin{aligned} MinI = & 2444469.4999(\pm 0.0016) + 0.40598942(\pm 0.00000020) \times E \\ & - 2.24(\pm 0.10) \times 10^{-10} E^2 + 0.0112(\pm 0.0012) \sin(0^\circ.0106E - 31^\circ.5). \end{aligned} \quad (9)$$

The resultant O-C diagrams are also shown in Figure 4 and 5, as the solid line represents. That fitting gives a long-term decrease ($dP/dt = -4.03 \times 10^{-7}$ days/year) with a clear period oscillation ($A_3 = 0^d.0112$, $T_3 = 37.75$ years). The residuals were plotted in Figure 6. Comparing with the Figure 3, the residuals in this graph is purely good.

4. Photometric solutions

Because of its unusual period changes, spectroscopic observation have been done by Lu (1991) and Rainger et al. (1992). Ten years later, Kim et al. (2003) gave a comprehensive investigation, including the period changes, the photometric and spectroscopic parameters. According to their studies, the spectroscopic mass ratio is more likely $q = 3.25$ which we adopted in our photometric solution. Of course, as a standard process, to check this value, a q-search method with the 2003 version of the W-D program (Wilson & Devinney, 1971, Wilson, 1990, 1994, Wilson & Van Hamme, 2003) was used (Figure 7). We fixed q to 0.3, 0.4, 0.5, and so on, as figure 7 shows. It can be seen that the best value is between $q = 3.1$ and $q = 3.7$, which is acceptable in the range of the errors.

During the solution, the temperature of star 1 (star eclipsed at primary light minimum) was fixed at $T_1 = 5860$ K, the same value used by Kim et al. (2003). The bolometric

albedo $A_1 = A_2 = 0.5$ (Rucinski 1969) and the values of the gravity-darkening coefficient $g_1 = g_2 = 0.32$ (Lucy 1967) were used, which correspond to the common convective envelope of both components. According to Claret & Gimenez (1990), a limb-darkening coefficient of 0.491 in R was used. We adjusted the orbital inclination i ; the mean temperature of star 2, T_2 ; the monochromatic luminosity of star 1, L_{1R} and the dimensionless potential of star 1 ($\Omega_1 = \Omega_2$, mode 3 for contact configuration). A small O’Connell effect of the system can not be ignored. As SS Ari’s spectral type is in the interval F8V-G2V, a sun-like star, it seemed that the probability of appearance of starspots on the surface of the star is high. In fact, as many researchers have done before (e.g., Liu et al. 1993, Kim et al. 2003), which were an agreement now, we add a spot on the star 2 that was subsequently proved as the cooler more massive primary component. The photometric solutions are listed in Table 3 and the theoretical light curves computed with those photometric elements are plotted in Figure 8. In order to get a image of the binary and its surface spots in our mind, the geometrical structure of SS Ari is displayed in Figure 9. For comparing, the results of previous studies are listed in Table 4.

5. Discussions and conclusions

The orbital period was revised as 0.40599166 days by using 115 photoelectric and CCD timings of SS Ari listed in Table 2. This system is a shallow contact W-Type binary with $q = 3.25$, $f = 9.4\%$. These suggest that the system may be a marginal contact binary. The temperatures of the two components are different (372K), that phenomena is very common in marginal contact binaries. Marginal contact binaries, whose fill factor are very small, ($f \leq 10\%$) are indicators of the time-scale that a binary will spend to evolve into the contact stage. For example, II CMa (Liu et al. 2008), V803 Aql (Samec et al. 1993), FG Sct (Bradstreet 1985), RW PsA (Lucy & Wilson 1979), XZ Leo (Niarchos et al. 1994), S Ant (Russo et al. 1982).

5.1. Model I

As shown in Figures 2, 3 and 4, both of the primary and the secondary times of light minimum follow the same general trend of O-C variation indicating that the weak O-C oscillation can barely be explained as apsidal motion. The alternate period change of a close binary containing at least one solar-type component at least can be interpreted by the mechanism of magnetic activity (e.g., Applegate 1992, Lanza et al. 1998). However, for contact binary stars, we scarcely know whether this mechanism can work or how it

might work; the common convective envelope covers the truth. We deduce that the period oscillation may be caused by the light-time effect of a tertiary component. The trend of O-C variation is very clearly, as we can see from Figure 2, though the data between E=0 and E=10000 show large scatters. In the previous section, by assuming an eccentric orbit and constant period model, a theoretical solution of the orbit for the assumed tertiary star was calculated. By using this equation:

$$f(m) = \frac{4\pi^2}{GT_3^2} \times (a'_{12} \sin i')^3, \quad (10)$$

(where $a'_{12} \sin i' = A_3 / \cos w' \times c / (1 \pm e')$; c is the speed of light; e' is the eccentricity of the third body's orbit; w' is the longitude of periastron of the third body's orbit), the mass function for the tertiary component is computed. Then, with the following equation:

$$f(m) = \frac{(M_3 \sin i')^3}{(M_1 + M_2 + M_3)^2}, \quad (11)$$

taking in the physical parameters $M_1 = 1.3M_\odot$, $M_2 = 0.4M_\odot$ (Kim et al. 2003), the masses and the orbital radii of the third companion are computed. The values for several different orbital inclinations (i') are shown in Table 5. If the tertiary companion is coplanar to the eclipsing pair (i.e., with the same inclination as the eclipsing binary), its mass should be $m_3 = 1.73M_\odot$. This mass is big enough to be detected. However, several spectral observations did not find a third body any more. Why is so massive star difficult to be seen? The possibility is that it is a compact star, what consistent in the results given by Kim et al. (2003). If the compact object really exists, the most point is how can the third body evolve into a compact star leaving a main sequence centric binary system? The interpretation by capture theory is more reasonable than by forming at same time and evolving together.

5.2. Model II

However, from Eq.9, we were amazed at the very small mass of the third body (About $0.278M_\odot$, see Table 5). We think that this model is a much better interpretation of the unseen tertiary. All doubts mentioned above will be swept away as the unseen component is a small mass star (e.g., the third body should be a cool dwarf star which is unseen spectroscopically and contributes extremely low light to the total system in optical band). This fitting gives its long-term decrease with a change rate $dP/dt = -4.03 \times 10^{-7}$ days/year, which may be due to a conservative mass transfer from the more massive component to the less massive one. Then with the accepted absolute parameters, the well-known equation (Tout & Hall 1991),

$$\frac{\dot{P}}{P} = 3 \frac{\dot{M}_2}{M_2} \left(\frac{M_2}{M_1} - 1 \right), \quad (12)$$

the mass transfer rate is estimated to be $dM_2/dt = -1.91 \times 10^{-7} M_\odot/\text{year}$ where minus expresses the more massive component M_2 losing mass. The timescale of mass transfer is $\tau \sim M_2/\dot{M}_2 \sim 6.80 \times 10^6$ years. This number is half of the thermal time scale of the massive component. On the other hand, both components of SS Ari are solar-type stars (spectral type in the interval F8V-G2V) but they rotate about 100 times as fast as the Sun, indicating a high degree of magnetic activity from the spin-up of the components. The asymmetry and variation of the light curve are indicators of spot activity. The long term decrease of the orbital period can be explained as the results of an enhanced stellar wind and AML. It is possible that the cyclic period change is caused by magnetic activity-driven variations in the quadrupole momentum of the solar-type components (e.g., Applegate 1992, Lanza et al. 1998).

Qian (2001a, b, 2003a) has shown the long-term period variation of contact binary stars may correlate with the mass of the primary component (M_1), the mass ratio of the system (q). His statistic critical mass ratio q is 0.4. When $q > 0.4$, the secular period increases; contrary, $q < 0.4$, the secular period decreases. The secular period decrease of SS Ari is consistent with this conclusion. To interpret the secular period changes of contact binary stars, Qian (2001a, b, 2003a) has proposed an evolutionary scenario of eclipsing binary stars. According to this scenario, the evolution of a contact binary may be the combination of the thermal relaxation oscillation (TRO) and the variable angular momentum loss (AML) via the change of depth of contact. Systems (e.g., V417 Aql, see Qian 2003b) with a secular decreasing period are on the AML-controlled stage, while those (e.g., CE Leo, see Qian 2002) showing an increasing period are on the TRO-controlled stage. The long-term period decrease of SS Ari may suggest that it is on the AML-controlled stage of this evolutionary scheme.

This work is partly supported by Yunnan Natural Science Foundation Foundation (No.2005A0059M), Chinese Natural Science Foundation (No.10573032, No.10433030, No.10573013 and No.10778707), The Ministry of Science and Technology of the Peoples Republic of China through grant 2007CB815406 and The Chinese Academy of Sciences grant No.O8ZKY11001 and No.O8AXB51001. New observations of the system were obtained with the 60cm telescope at Yunnan Observatory. Thanks to the anonymous referee who has given us very constructive comments and cordial suggestions, which helped us to improve the paper greatly.

REFERENCES

Agerer, F., Hubscher, J., 2003, IBVS 5484, 1A

- Applegate, J. H., 1992, ApJ 385, 621
- Barden, S. C., 1987, ApJ 317, 333
- Biro, I. B., Borkovits, T., Hegedus, T., Kiss, Z. T., Kovacs, T., Lampens, P., Regaly, Zs., Robertson, C. W., van Cauteren, P., 2007, IBVS 5760, 1N
- Borkovits, T., Biro, I. B., Hegedus, T., Csizmadia, S., Szabados, L., Pal, A., Posztobanyi, K., Konyves, V., Kospal, A., Csak, B., Meszaros, S., 2003, IBVS 5434, 1B
- Bradstreet, D. H., 1985, ApJS 58, 413B
- Braune, W., 1970, IBVS 440
- Braune, W., Hubscher, J. & Mundry, E., 1972, AN 294, 123
- Braune, W. & Mundry, E., 1973, AN 294, 225
- Braune, W., Hubscher, J. & Mundry, E., 1977, AN 298, 121
- Braune, W., Hubscher, J. & Mundry, E., 1979, AN 300, 165
- Braune, W., Hubscher, J. & Mundry, E., 1981, AN 302, 53
- Braune, W., Hubscher, J. & Mundry, E., 1983, BAV Bull. No. 36
- Claret, A., and Gimenez, A., 1990, A&A 230, 412
- Demircan, O. and Selam S. O., 1993, A&A 267, 107
- Diethelm, R., 1992, BBSAG Bull. No. 99
- Diethelm, R. & Blattler, E., 1998, BBSAG Bull. No. 116
- Faulkner, D. R., 1986, PASP 98, 690
- Hanzl, D. R., 1990, IBVS 3423
- Hegedus, T., Biro, I. B., Borkovits, T., and Paragi, 1996, IBVS 4340
- Hoffmeister, C., 1934, Astron. Nachr. 253, 195
- Hubscher, J., 2005, IBVS 5643, 1H
- Hubscher, J. & Mundry, E., 1984, BAV Bull. No. 38
- Hubscher, J., Lichtenknecker, D. & Meyer, J., 1986, BAV Bull. No. 43
- Hubscher, J. & Lichtenknecker, D., 1988, BAV Bull. No. 50
- Hubscher, J., Agerer, F. & Wunder, E., 1992, BAV Mitt. No. 60
- Hubscher, J., Agerer, F. & Wunder, E., 1993, BAV Mitt. No. 62
- Hubscher, J., Paschke, A., Walter, F., 2005, IBVS 5657, 1H
- Hubscher, J., Paschke, A., Walter, F., 2006, IBVS 5731, 1H

- Hubscher, J., & Walter, F., 2007, IBVS 5761, 1H
- Huth, H., 1964, Mitt. Sonnebreg 2, 112
- Isles, J. E., 1985a, BAA Var. Star Section Circ. No. 60
- Isles, J. E., 1985b, BAA Var. Star Section Circ. No. 61
- Isles, J. E., 1986, BAA Var. Star Section Circ. No. 63
- Isles, J. E., 1988, BAA Var. Star Section Circ. No. 66
- Kaluzny, J. and Pojmanski, G., 1984a, IBVS 2564
- Kaluzny, J. and Pojmanski, G., 1984b, Acta. Astron. 34, 445
- Kim, C.-H., Han, W., Yoon, J. H. & Nha, I.-S., 1997, J. Astron. Space Sci. 14, 44
- Kim, Chun-Hwey, Lee, Jae-Woo, Kim, Seung-Lee, Han, Wonyong, Koch, Robert H., 2003, AJ, 125, 322K
- Kopal, D., 1978, Dynamics of Close binary Systems, D. Reidel. Co.
- Krajci, Tom., 2005, IBVS 5592, 1K
- Kramer, E. H., 1948, Astr. Astrophys. 128, 84
- Kurpinska-Winiarska, M. and Zakrzewski, B., 1990, IBVS 3485
- Lanza, A. F., Rodonò, M., and Rosner, R., 1998, MNRAS 296, 893
- Liu Liang, Qian Sheng-Bang, Zhu Li-Ying, He Jia-Jia, Yuan Jin-Zhao, Dai Zhi-Bin, Liao Wen-Ping and Zhang Jia, 2008, PASJ 60, 3
- Liu Qingyao, Yang Yulan, Gu Shenghong, and Wang Bi, 1996, A&As 101, 253
- Locher, K., 1978, BBSAG Bull. No. 39
- Locher, K., 1980a, BBSAG Bull. No. 46
- Locher, K., 1980b, BBSAG Bull. No. 51
- Locher, K., 1981a, BBSAG Bull. No. 52
- Locher, K., 1981b, BBSAG Bull. No. 53
- Locher, K., 1981c, BBSAG Bull. No. 56
- Locher, K., 1981d, BBSAG Bull. No. 57
- Locher, K., 1982a, BBSAG Bull. No. 58
- Locher, K., 1982b, BBSAG Bull. No. 62
- Locher, K., 1983a, BBSAG Bull. No. 64
- Locher, K., 1983b, BBSAG Bull. No. 65

- Locher, K., 1983c, BBSAG Bull. No. 69
- Locher, K., 1984a, BBSAG Bull. No. 70
- Locher, K., 1984b, BBSAG Bull. No. 71
- Locher, K., 1984c, BBSAG Bull. No. 74
- Locher, K., 1985, BBSAG Bull. No. 78
- Locher, K., 1986, BBSAG Bull. No. 79
- Locher, K., 1987a, BBSAG Bull. No. 82
- Locher, K., 1987b, BBSAG Bull. No. 83
- Locher, K., 1988a, BBSAG Bull. No. 86
- Locher, K., 1988b, BBSAG Bull. No. 87
- Locher, K., 1988c, BBSAG Bull. No. 88
- Locher, K., 1989a, BBSAG Bull. No. 90
- Locher, K., 1989b, BBSAG Bull. No. 91
- Lucy, L. B., Wilson, R. E., 1979, ApJ 231, 502L
- Lu Wenxian, 1991, AJ 102, 262L
- Muyesseroglu, Z., Burol, B., and Selam, S. O., 1996, IBVS 4380
- Nelson, R. H., 2001, IBVS 5040
- Nelson, R. H., 2007, IBVS 5760
- Niarchos, P. G., Hoffmann, M., Duerbeck, H. W., 1994, A&A 292, 494N
- Odynskaya, O. K., 1949, Perem. Zvezdy 6, 316
- Ogloza, W., 1995, IBVS 4263
- Ogloza, W., 1997, IBVS 4534
- Pohl, E., Hamzaoglu, E., Gudur, N. & Ibanoglu, C., 1983, IBVS 2385
- Pohl, E., Tunca, Z., Gulmen, O. & Evren, S., 1985, IBVS 2793
- Pohl, E., Akan, M. C., Sezer, C. & Gudur, N., 1987, IBVS 3087
- Pribulla, T., Vanko, M., Parimucha, S. & Chochol, D., 2001, IBVS 5056
- Pribulla, T., Vanko, M., Parimucha, S., Chochol, D., 2002, IBVS 5341, 1P
- Qian, S.-B., 2001a, MNRAS 328, 635
- Qian, S.-B., 2001b, MNRAS 328, 914

- Qian, S.-B., 2002, *A&A* 384, 908
- Qian, S.-B., 2003a, *MNRAS* 342, 1260
- Qian, S.-B., 2003b, *A&A* 400, 649
- Qian, S.-B., He, J.-J., Xiang, F.-Y., Ding, X.-Y., Soonthornthum, B., 2005, *AJ* 129, 1686Q
- Qian, S.-B., Zhu, L.-Y., Boonruksar, S., 2006, *NewA* 11 503Q
- Qian, S.-B., Xiang, F.-Y., Zhu, L.-Y., Dai, Z.-B., He, J.-J., Yuan, J.-Z., 2007, *AJ* 133, 357Q
- Rainger, P. P., Bell, S. A., Hilditch, R. W., 1992, *MNRAS*, 254, 568R
- Russo, G., Sollazzo, C., Maceroni, C., Milano, L., 1982, *A&AS* 47, 211R
- Rucinski, S. M., 1969, *A&A* 19, 245
- Samec, Ronald G., Su, Wen, Dewitt, Jason R., 1993, *PASP* 105, 1441S
- Selam, S. O., Gural, B., and Muyesseroglu, Z., 1999, *IBVS* 4670
- Tout, C. A. & Hall, D. S., 1991, *MNRAS* 253, 9T
- Wilson R. E. 1990, *ApJ* 356, 613
- Wilson R. E. 1994, *PASP* 106, 921
- Wilson R. E. & Devinney, E. J., 1971, *ApJ* 166, 605
- Wilson R. E. & Van Hamme, W., 2003, *Computing Binary Stars Observables*, the 4th edition of the W-D programe.
- Voges, W., Aschenbach, B., Boller, Th., Brauninger, H., Briel, U., Burkert, W., Dennerl, K., Englhauser, J., Gruber, R., Haberl, F., and 8 coauthors, 2000, *yCat* 33560445B
- Zhukov, G. V., 1975, *Astron. Tsirk.* No. 888, 7

Table 1: The original data of SS Ari in R band observed by 60cm telescope at Yunnan Observatory. Hel. JD 2454400+

Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm
47.9884	1.158	48.0343	0.823	48.0799	0.713	48.1287	0.798	48.1754	1.150	48.2281	0.899	48.2861	0.674
47.9897	1.151	48.0348	0.824	48.0804	0.705	48.1292	0.797	48.1760	1.157	48.2290	0.884	48.2872	0.665
47.9902	1.155	48.0354	0.819	48.0810	0.704	48.1298	0.794	48.1766	1.149	48.2295	0.874	48.2880	0.693
47.9913	1.146	48.0360	0.816	48.0816	0.705	48.1303	0.795	48.1772	1.162	48.2302	0.882	48.2888	0.690
47.9919	1.148	48.0366	0.809	48.0821	0.699	48.1311	0.809	48.1777	1.162	48.2313	0.877	48.2895	0.666
47.9924	1.143	48.0372	0.814	48.0828	0.689	48.1318	0.807	48.1783	1.173	48.2320	0.856	48.2901	0.671
47.9929	1.141	48.0378	0.805	48.0834	0.692	48.1323	0.809	48.1789	1.175	48.2331	0.855	48.2907	0.668
47.9934	1.144	48.0383	0.800	48.0840	0.694	48.1329	0.804	48.1796	1.179	48.2341	0.847	48.2914	0.665
47.9940	1.136	48.0388	0.800	48.0845	0.694	48.1335	0.814	48.1802	1.196	48.2347	0.848	48.2944	0.681
47.9944	1.132	48.0394	0.796	48.0851	0.692	48.1344	0.818	48.1818	1.186	48.2353	0.838	48.2956	0.686
47.9949	1.138	48.0399	0.801	48.0856	0.696	48.1349	0.819	48.1829	1.192	48.2359	0.832	48.2964	0.670
47.9954	1.129	48.0406	0.801	48.0862	0.688	48.1354	0.824	48.1834	1.188	48.2364	0.836	48.2970	0.666
47.9960	1.127	48.0411	0.785	48.0868	0.695	48.1359	0.818	48.1840	1.202	48.2371	0.837	48.2978	0.674
47.9965	1.124	48.0417	0.794	48.0874	0.694	48.1365	0.823	48.1846	1.202	48.2377	0.839	48.2984	0.696
47.9971	1.120	48.0422	0.787	48.0880	0.696	48.1370	0.837	48.1852	1.202	48.2384	0.819	48.2990	0.687
47.9976	1.125	48.0428	0.783	48.0886	0.702	48.1375	0.827	48.1857	1.215	48.2395	0.821	48.2996	0.709
47.9981	1.119	48.0433	0.788	48.0892	0.699	48.1380	0.827	48.1863	1.206	48.2400	0.819	48.3002	0.682
47.9990	1.109	48.0439	0.779	48.0897	0.697	48.1386	0.837	48.1869	1.214	48.2406	0.808	48.3007	0.681
47.9995	1.106	48.0444	0.780	48.0902	0.699	48.1391	0.838	48.1874	1.215	48.2412	0.806	48.3015	0.688
48.0000	1.094	48.0450	0.774	48.0908	0.700	48.1398	0.847	48.1880	1.214	48.2418	0.797	48.3027	0.680
48.0006	1.095	48.0455	0.781	48.0914	0.703	48.1404	0.851	48.1885	1.207	48.2425	0.792	48.3039	0.687
48.0011	1.092	48.0461	0.777	48.0919	0.692	48.1409	0.851	48.1890	1.215	48.2433	0.801	48.3050	0.670
48.0016	1.084	48.0467	0.782	48.0925	0.697	48.1415	0.852	48.1896	1.211	48.2440	0.797	48.3059	0.704
48.0022	1.081	48.0472	0.776	48.0930	0.712	48.1420	0.856	48.1903	1.226	48.2450	0.790	48.3066	0.667
48.0027	1.074	48.0478	0.772	48.0936	0.701	48.1426	0.859	48.1909	1.205	48.2456	0.788	48.3072	0.662
48.0032	1.072	48.0484	0.776	48.0942	0.709	48.1432	0.857	48.1915	1.210	48.2462	0.789	48.3079	0.678
48.0038	1.069	48.0490	0.777	48.0947	0.698	48.1437	0.870	48.1921	1.209	48.2468	0.790	48.3086	0.699
48.0043	1.058	48.0495	0.771	48.0953	0.709	48.1442	0.875	48.1926	1.178	48.2476	0.778	48.3094	0.711
48.0049	1.064	48.0501	0.766	48.0959	0.706	48.1448	0.867	48.1932	1.208	48.2484	0.774	48.3100	0.698
48.0054	1.059	48.0506	0.763	48.0964	0.705	48.1454	0.878	48.1938	1.201	48.2489	0.778	48.3106	0.689
48.0060	1.055	48.0512	0.769	48.0970	0.711	48.1460	0.882	48.1943	1.208	48.2495	0.780	48.3112	0.671
48.0065	1.055	48.0520	0.761	48.0975	0.707	48.1465	0.887	48.1949	1.198	48.2501	0.766	48.3119	0.719
48.0070	1.038	48.0526	0.762	48.0981	0.703	48.1470	0.883	48.1955	1.192	48.2509	0.777	48.3125	0.689
48.0077	1.015	48.0532	0.757	48.0987	0.715	48.1476	0.895	48.1960	1.178	48.2515	0.764	48.3132	0.685
48.0082	1.017	48.0537	0.756	48.0992	0.710	48.1482	0.902	48.1966	1.191	48.2521	0.757	48.3138	0.655
48.0088	1.002	48.0544	0.759	48.0998	0.712	48.1487	0.907	48.1972	1.190	48.2526	0.715	48.3146	0.682
48.0093	1.006	48.0550	0.758	48.1003	0.716	48.1494	0.911	48.1978	1.191	48.2532	0.716	48.3156	0.699
48.0099	0.999	48.0555	0.758	48.1010	0.723	48.1499	0.927	48.1984	1.190	48.2537	0.742	48.3163	0.703
48.0105	0.994	48.0568	0.738	48.1016	0.719	48.1505	0.911	48.1990	1.198	48.2546	0.749	48.3170	0.713
48.0110	0.993	48.0574	0.734	48.1021	0.721	48.1511	0.926	48.1996	1.165	48.2552	0.750	48.3178	0.684
48.0115	0.981	48.0579	0.737	48.1026	0.720	48.1516	0.927	48.2001	1.163	48.2559	0.729	48.3183	0.698
48.0121	0.982	48.0584	0.735	48.1032	0.721	48.1521	0.922	48.2007	1.156	48.2565	0.737	48.3190	0.728
48.0126	0.971	48.0590	0.735	48.1038	0.721	48.1539	0.944	48.2033	1.130	48.2574	0.732	48.3195	0.692
48.0132	0.966	48.0596	0.733	48.1043	0.730	48.1547	0.938	48.2039	1.131	48.2583	0.746	48.3208	0.701
48.0137	0.969	48.0601	0.730	48.1048	0.727	48.1553	0.945	48.2051	1.115	48.2590	0.747	48.3215	0.737
48.0142	0.972	48.0607	0.726	48.1061	0.727	48.1558	0.952	48.2057	1.110	48.2602	0.757	48.3221	0.719
48.0147	0.954	48.0612	0.735	48.1066	0.727	48.1563	0.951	48.2065	1.103	48.2610	0.724	48.3226	0.714
48.0152	0.957	48.0617	0.726	48.1078	0.732	48.1569	0.959	48.2071	1.104	48.2615	0.743	48.3233	0.693
48.0158	0.949	48.0623	0.728	48.1084	0.738	48.1574	0.966	48.2078	1.096	48.2622	0.755	48.3241	0.746
48.0165	0.943	48.0628	0.725	48.1089	0.734	48.1579	0.971	48.2084	1.098	48.2628	0.748	48.3246	0.751
48.0170	0.940	48.0633	0.724	48.1095	0.740	48.1586	0.976	48.2090	1.080	48.2636	0.742	48.3252	0.701
48.0175	0.937	48.0639	0.724	48.1101	0.740	48.1591	0.987	48.2095	1.075	48.2643	0.731	48.3257	0.731
48.0181	0.937	48.0644	0.724	48.1106	0.739	48.1598	0.997	48.2101	1.081	48.2656	0.718	48.3262	0.732
48.0187	0.924	48.0650	0.720	48.1111	0.746	48.1603	0.993	48.2106	1.072	48.2669	0.722	48.3268	0.711
48.0192	0.922	48.0656	0.717	48.1117	0.751	48.1610	0.998	48.2112	1.061	48.2675	0.707	48.3274	0.711
48.0198	0.923	48.0661	0.717	48.1122	0.749	48.1616	1.004	48.2118	1.060	48.2681	0.713	48.3280	0.721
48.0203	0.913	48.0667	0.714	48.1128	0.749	48.1622	1.016	48.2124	1.055	48.2686	0.717	48.3286	0.770
48.0209	0.914	48.0672	0.705	48.1134	0.766	48.1627	1.026	48.2129	1.042	48.2705	0.709	48.3298	0.756
48.0214	0.913	48.0678	0.717	48.1139	0.749	48.1634	1.030	48.2140	1.021	48.2714	0.704	48.3304	0.778
48.0220	0.899	48.0683	0.710	48.1145	0.758	48.1639	1.038	48.2145	1.021	48.2721	0.705	48.3311	0.713
48.0226	0.898	48.0688	0.716	48.1150	0.757	48.1644	1.035	48.2152	1.015	48.2727	0.709	48.3316	0.761
48.0231	0.901	48.0694	0.717	48.1155	0.758	48.1649	1.044	48.2158	1.009	48.2733	0.704	48.3325	0.732
48.0236	0.895	48.0700	0.708	48.1161	0.754	48.1655	1.052	48.2165	1.001	48.2739	0.704	48.3333	0.800
48.0242	0.895	48.0705	0.713	48.1166	0.758	48.1661	1.052	48.2172	0.990	48.2745	0.713	48.3342	0.797
48.0248	0.891	48.0710	0.706	48.1172	0.760	48.1666	1.054	48.2180	0.991	48.2751	0.710	48.3347	0.714
48.0253	0.881	48.0716	0.715	48.1177	0.765	48.1672	1.064	48.2185	0.982	48.2758	0.707	48.3354	0.821
48.0258	0.882	48.0721	0.716	48.1182	0.762	48.1677	1.073	48.2190	0.977	48.2766	0.697	48.3359	0.753
48.0264	0.875	48.0727	0.718	48.1187	0.749	48.1683	1.080	48.2196	0.973	48.2771	0.689	48.3365	0.790
48.0269	0.870	48.0733	0.708	48.1216	0.771	48.1689	1.086	48.2203	0.960	48.2779	0.688	48.3371	0.723
48.0283	0.854	48.0738	0.709	48.1222	0.769	48.1694	1.092	48.2208	0.962	48.2785	0.690	48.3376	0.700
48.0289	0.844	48.0744	0.709	48.1223	0.769	48.1699	1.092	48.2215	0.949	48.2791	0.691	48.3391	0.741
48.0294	0.845	48.0749	0.713	48.1236	0.779	48.1705	1.101	48.2220	0.949	48.2798	0.675	48.3397	0.772
48.0299	0.847	48.0754	0.708	48.1242	0.781	48.1711	1.106	48.2227	0.936	48.2805	0.696	48.3416	0.697
48.0305	0.842	48.0761	0.704	48.1247	0.785	48.1716	1.118	48.2233	0.939	48.2812	0.691	48.3422	0.885
48.0311	0.838	48.0766	0.705	48.1253	0.781	48.1722	1.109	48.2239	0.932	48.2819	0.678	48.3428	0.753
48.0316	0.831	48.0772	0.701	48.1259	0.790	48.1727	1.114	48.2246	0.917	48.2825	0.688	48.3435	0.766
48.0321													

Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm	Hel.JD	Δm
48.3472	0.788	49.0971	0.690	49.1242	0.698	49.1514	0.776	49.1788	0.955	49.2089	1.126
48.3477	0.728	49.0974	0.672	49.1245	0.704	49.1517	0.778	49.1791	0.967	49.2092	1.134
48.3484	0.873	49.0978	0.681	49.1248	0.702	49.1521	0.795	49.1794	0.958	49.2095	1.132
48.3491	0.806	49.0982	0.686	49.1251	0.711	49.1524	0.773	49.1797	0.982	49.2098	1.122
49.0706	0.732	49.0987	0.688	49.1255	0.717	49.1527	0.797	49.1800	0.967	49.2102	1.119
49.0710	0.734	49.0991	0.673	49.1258	0.704	49.1531	0.781	49.1804	0.969	49.2105	1.123
49.0714	0.736	49.0994	0.683	49.1262	0.706	49.1534	0.778	49.1807	0.971	49.2108	1.144
49.0718	0.737	49.0997	0.670	49.1265	0.711	49.1537	0.799	49.1811	0.983	49.2112	1.119
49.0721	0.738	49.1000	0.690	49.1268	0.716	49.1540	0.793	49.1814	0.981	49.2115	1.124
49.0724	0.732	49.1004	0.680	49.1271	0.710	49.1544	0.795	49.1817	0.975	49.2118	1.104
49.0728	0.733	49.1007	0.681	49.1274	0.720	49.1547	0.802	49.1820	0.994	49.2122	1.132
49.0731	0.730	49.1010	0.692	49.1277	0.725	49.1550	0.803	49.1824	1.003	49.2126	1.129
49.0735	0.734	49.1013	0.688	49.1280	0.713	49.1553	0.800	49.1827	1.003	49.2130	1.122
49.0738	0.738	49.1017	0.695	49.1283	0.730	49.1557	0.791	49.1832	1.003	49.2133	1.102
49.0741	0.729	49.1020	0.683	49.1287	0.711	49.1560	0.796	49.1835	0.996	49.2137	1.120
49.0744	0.733	49.1024	0.687	49.1290	0.719	49.1563	0.806	49.1839	1.007	49.2140	1.094
49.0748	0.746	49.1027	0.699	49.1293	0.722	49.1566	0.812	49.1842	1.004	49.2144	1.108
49.0751	0.744	49.1031	0.700	49.1296	0.714	49.1569	0.819	49.1845	1.025	49.2147	1.112
49.0754	0.724	49.1034	0.688	49.1301	0.714	49.1573	0.818	49.1848	1.015	49.2150	1.075
49.0757	0.730	49.1037	0.683	49.1304	0.718	49.1576	0.809	49.1851	1.018	49.2153	1.086
49.0760	0.722	49.1041	0.689	49.1308	0.711	49.1579	0.808	49.1855	1.021	49.2156	1.108
49.0763	0.728	49.1045	0.693	49.1312	0.721	49.1582	0.790	49.1858	1.028	49.2159	1.105
49.0766	0.723	49.1048	0.689	49.1315	0.727	49.1586	0.795	49.1861	1.029	49.2163	1.117
49.0769	0.713	49.1051	0.680	49.1319	0.730	49.1590	0.798	49.1865	1.037	49.2167	1.061
49.0773	0.716	49.1055	0.699	49.1322	0.721	49.1594	0.775	49.1868	1.027	49.2171	1.090
49.0776	0.725	49.1058	0.692	49.1326	0.718	49.1597	0.772	49.1871	1.024	49.2174	1.091
49.0780	0.716	49.1061	0.694	49.1329	0.719	49.1600	0.784	49.1874	1.051	49.2177	1.072
49.0783	0.711	49.1065	0.694	49.1332	0.713	49.1603	0.800	49.1878	1.052	49.2181	1.096
49.0786	0.718	49.1068	0.690	49.1335	0.721	49.1607	0.805	49.1881	1.059	49.2184	1.095
49.0789	0.710	49.1071	0.687	49.1338	0.735	49.1610	0.800	49.1885	1.065	49.2188	1.074
49.0792	0.718	49.1074	0.677	49.1341	0.729	49.1613	0.786	49.1888	1.052	49.2192	1.070
49.0795	0.714	49.1077	0.679	49.1345	0.723	49.1617	0.810	49.1892	1.074	49.2195	1.071
49.0799	0.720	49.1081	0.696	49.1348	0.731	49.1621	0.798	49.1895	1.060	49.2198	1.055
49.0802	0.720	49.1084	0.693	49.1351	0.734	49.1625	0.801	49.1899	1.070	49.2202	1.070
49.0806	0.723	49.1087	0.697	49.1354	0.751	49.1628	0.811	49.1905	1.075	49.2206	1.066
49.0809	0.715	49.1093	0.688	49.1357	0.729	49.1631	0.786	49.1908	1.086	49.2209	1.043
49.0812	0.719	49.1096	0.679	49.1361	0.742	49.1634	0.812	49.1912	1.089	49.2212	1.066
49.0815	0.710	49.1101	0.674	49.1364	0.733	49.1637	0.794	49.1915	1.058	49.2216	1.071
49.0819	0.706	49.1104	0.690	49.1367	0.732	49.1640	0.802	49.1918	1.072	49.2219	1.057
49.0822	0.710	49.1107	0.684	49.1370	0.740	49.1644	0.797	49.1923	1.090	49.2223	1.055
49.0825	0.715	49.1111	0.688	49.1373	0.745	49.1647	0.805	49.1956	1.108	49.2226	1.038
49.0828	0.705	49.1114	0.696	49.1376	0.726	49.1650	0.798	49.1959	1.122	49.2230	1.050
49.0831	0.704	49.1117	0.703	49.1380	0.745	49.1653	0.815	49.1963	1.112	49.2233	1.016
49.0835	0.712	49.1120	0.689	49.1383	0.746	49.1657	0.803	49.1966	1.132	49.2236	1.047
49.0840	0.721	49.1124	0.692	49.1387	0.747	49.1661	0.805	49.1969	1.125	49.2240	1.023
49.0844	0.711	49.1127	0.693	49.1390	0.744	49.1664	0.798	49.1973	1.133	49.2243	1.016
49.0848	0.717	49.1131	0.698	49.1393	0.747	49.1667	0.827	49.1976	1.121	49.2246	1.042
49.0851	0.711	49.1134	0.686	49.1396	0.738	49.1670	0.806	49.1979	1.122	49.2252	1.023
49.0854	0.716	49.1137	0.699	49.1399	0.743	49.1673	0.797	49.1983	1.127	49.2255	1.036
49.0858	0.707	49.1140	0.697	49.1403	0.759	49.1677	0.816	49.1986	1.115	49.2259	1.026
49.0861	0.711	49.1143	0.704	49.1406	0.750	49.1681	0.799	49.1989	1.118	49.2263	1.017
49.0865	0.709	49.1147	0.707	49.1409	0.750	49.1684	0.807	49.1992	1.136	49.2266	1.008
49.0868	0.695	49.1150	0.687	49.1412	0.744	49.1691	0.808	49.1995	1.101		
49.0872	0.706	49.1153	0.694	49.1415	0.753	49.1695	0.815	49.1999	1.153		
49.0876	0.708	49.1156	0.678	49.1419	0.738	49.1701	0.848	49.2002	1.128		
49.0879	0.706	49.1159	0.688	49.1422	0.758	49.1704	0.907	49.2006	1.138		
49.0882	0.702	49.1162	0.701	49.1425	0.750	49.1708	0.892	49.2009	1.156		
49.0890	0.707	49.1165	0.698	49.1428	0.769	49.1711	0.899	49.2012	1.133		
49.0897	0.694	49.1168	0.691	49.1431	0.751	49.1714	0.900	49.2016	1.141		
49.0901	0.705	49.1172	0.694	49.1435	0.752	49.1718	0.911	49.2019	1.123		
49.0904	0.687	49.1175	0.701	49.1438	0.757	49.1721	0.919	49.2023	1.145		
49.0907	0.707	49.1178	0.693	49.1441	0.749	49.1725	0.917	49.2026	1.145		
49.0911	0.695	49.1181	0.707	49.1444	0.757	49.1728	0.935	49.2030	1.159		
49.0914	0.687	49.1184	0.696	49.1448	0.783	49.1731	0.930	49.2033	1.146		
49.0917	0.690	49.1188	0.700	49.1451	0.757	49.1734	0.919	49.2036	1.145		
49.0921	0.687	49.1193	0.704	49.1455	0.761	49.1738	0.921	49.2039	1.145		
49.0924	0.694	49.1196	0.706	49.1459	0.761	49.1741	0.935	49.2042	1.143		
49.0927	0.689	49.1199	0.691	49.1462	0.775	49.1744	0.940	49.2046	1.136		
49.0931	0.689	49.1203	0.685	49.1465	0.756	49.1748	0.926	49.2049	1.143		
49.0934	0.681	49.1206	0.699	49.1468	0.798	49.1751	0.934	49.2052	1.162		
49.0937	0.691	49.1210	0.703	49.1471	0.780	49.1755	0.940	49.2056	1.144		
49.0940	0.674	49.1213	0.692	49.1475	0.776	49.1758	0.940	49.2059	1.117		
49.0944	0.687	49.1216	0.690	49.1478	0.779	49.1761	0.959	49.2062	1.147		
49.0947	0.696	49.1219	0.688	49.1481	0.765	49.1764	0.937	49.2065	1.141		
49.0951	0.679	49.1223	0.701	49.1495	0.779	49.1768	0.943	49.2069	1.153		
49.0954	0.693	49.1226	0.699	49.1498	0.746	49.1771	0.944	49.2072	1.130		
49.0957	0.692	49.1229	0.703	49.1501	0.785	49.1774	0.939	49.2075	1.124		
49.0960	0.679	49.1232	0.708	49.1504	0.777	49.1777	0.950	49.2079	1.177		
49.0963	0.685	49.1235	0.699	49.1508	0.782	49.1781	0.953	49.2082	1.104		
49.0966	0.674	49.1239	0.701	49.1511	0.783	49.1784	0.951	49.2085	1.134		

Table 2. Times of light minimum of SS Ari.

Hel.JD	Type	Method	E	($O - C$) Eq.2	($O - C$) Eq.7	Reference
2430948.3290	I	vis	-33304	-0.1915	-	Odynskaya (1949)
2432455.3470	I	vis	-29592	-0.1946	-	Kramer (1948)
2432455.5520	II	vis	-29591.5	-0.1926	-	Kramer (1948)
2432786.2290	I	vis	-28777	-0.1914	-	Kramer (1948)
2432786.4300	II	vis	-28776.5	-0.1934	-	Kramer (1948)
2435721.5450	I	pg	-21547	-0.1563	-	Huth (1964)
2436075.5810	I	pg	-20675	-0.1404	-	Huth (1964)
2439028.3870	I	vis	-13402	-0.0726	-	Braune (1970)
2439029.6090	I	vis	-13399	-0.0686	-	Braune (1970)
2439040.5713	I	pe	-13372	-0.0679	0.0129	Kaluzny & Pojmanski (1984)
2439053.3620	II	vis	-13340.5	-0.0658	-	Braune (1970)
2439055.3940	II	vis	-13335.5	-0.0637	-	Braune (1970)
2439068.3960	II	vis	-13303.5	-0.0533	-	Braune (1970)
2439184.3010	I	vis	-13018	-0.0574	-	Braune (1970)
2439389.5261	II	pe	-12512.5	-0.0584	0.0179	Kaluzny & Pojmanski (1984b)
2439389.5350	II	vis	-12512.5	-0.0495	-	Braune (1970)
2439391.5552	II	pe	-12507.5	-0.0592	0.0170	Kaluzny & Pojmanski (1984b)
2439403.5360	I	vis	-12478	-0.0550	-	Braune (1970)
2439407.5930	I	vis	-12468	-0.0578	-	Braune (1970)
2439776.4355	II	vis	-11559.5	-0.0539	-	Braune (1970)
2440065.5180	II	vis	-10847.5	-0.0336	-	Braune (1970)
2441249.3920	II	vis	-7931.5	-0.0156	-	Braune et al. (1972)
2441576.4220	I	vis	-7126	-0.0076	-	Braune & Mundry (1973)
2441682.3760	I	vis	-6865	-0.0160	-	Braune & Mundry (1973)
2441947.4934	I	pe	-6212	-0.0077	0.0347	Zhukov (1975)
2441951.5542	I	pe	-6202	-0.0067	0.0356	Zhukov (1975)
2441960.4865	I	pe	-6180	-0.0061	0.0361	Zhukov (1975)
2441975.5072	I	pe	-6143	-0.0069	0.0351	Zhukov (1975)
2442036.2053	II	pe	-5993.5	-0.0038	0.0374	Zhukov (1975)
2442037.4210	II	pe	-5990.5	-0.0060	0.0351	Zhukov (1975)
2442414.2010	II	vis	-5062.5	0.0187	-	Braune et al. (1977)
2442664.4740	I	vis	-4446	0.0012	-	Braune et al. (1979)
2442727.4086	I	pe	-4291	0.0079	0.0399	Kurpinska-Winiarska & Zakrzewski (1990)
2442758.2642	I	pe	-4215	0.0085	0.0402	Kurpinska-Winiarska & Zakrzewski (1990)
2442759.2791	II	pe	-4212.5	0.0085	0.0401	Kurpinska-Winiarska & Zakrzewski (1990)
2442840.2650	I	vis	-4013	0.0001	-	Braune et al. (1979)
2443014.4590	I	vis	-3584	0.0260	-	Braune et al. (1979)

Table 2—Continued

Hel.JD	Type	Method	E	($O - C$) Eq.2	($O - C$) Eq.7	Reference
2443833.3320	I	vis	-1567	0.0246	-	Braune et al. (1981)
2443455.3481	I	pe	-2498	0.0140	0.0364	Kurpinska-Winiarska & Zakrzewski (1990)
2443790.3000	I	vis	-1673	0.0272	-	Locher (1978)
2443795.5730	I	vis	-1660	0.0223	-	Locher (1978)
2443815.4581	I	pe	-1611	0.0141	0.0318	Kurpinska-Winiarska & Zakrzewski (1990)
2443927.3091	II	pe	-1335.5	0.0159	0.0321	Kurpinska-Winiarska & Zakrzewski (1990)
2444146.5426	II	pe	-795.5	0.0168	0.0301	Kurpinska-Winiarska & Zakrzewski (1990)
2444266.3180	II	vis	-500.5	0.0262	-	Locher (1980a)
2444469.5074	I	pe	0	0.0195	0.0285	Kurpinska-Winiarska & Zakrzewski (1990)
2444539.3260	I	vis	172	0.0085	-	Locher (1980b)
2444602.2450	I	vis	327	-0.0004	-	Locher (1981a)
2444605.2890	II	vis	334.5	-0.0013	-	Locher (1981a)
2444605.3104	II	pe	334.5	0.0201	0.0273	Kurpinska-Winiarska & Zakrzewski (1990)
2444629.2580	II	vis	393.5	0.0145	-	Locher (1981a)
2444635.3190	II	vis	408.5	-0.0143	-	Locher (1981a)
2444636.3420	I	vis	411	-0.0063	-	Locher (1981a)
2444642.2567	II	pe	425.5	0.0216	0.0284	Kurpinska-Winiarska & Zakrzewski (1990)
2444649.3240	I	vis	443	-0.0158	-	Locher (1981b)
2444659.2690	II	vis	467.5	-0.0175	-	Locher (1981b)
2444821.4560	I	vis	867	-0.0220	-	Locher (1981c)
2444823.5267	I	pe	872	0.0188	0.0231	Kurpinska-Winiarska & Zakrzewski (1990)
2444831.6060	I	vis	892	-0.0217	-	Locher (1981c)
2444879.3250	II	vis	1009.5	-0.0061	-	Locher (1981d)
2444883.3800	II	vis	1019.5	-0.0109	-	Locher (1981d)
2444911.4240	II	pe	1088.5	0.0200	0.0232	Locher (1981d)
2444917.2780	I	vis	1103	-0.0128	-	Locher (1981d)
2444919.3050	I	vis	1108	-0.0157	-	Locher (1981d)
2444926.2350	I	vis	1125	0.0125	-	Locher (1981d)
2444929.2690	II	vis	1132.5	0.0016	-	Locher (1981d)
2444985.2980	II	vis	1270.5	0.0045	-	Locher (1982a)
2445224.4580	II	vis	1859.5	0.0386	-	Locher (1982b)
2445238.4500	I	pg	1894	0.0241	-	Braune et al. (1983)
2445261.3848	II	pe	1950.5	0.0206	0.0192	Kaluzny & Pojmanski (1984b)
2445261.5860	I	pe	1951	0.0189	0.0174	Kaluzny & Pojmanski (1984b)
2445262.3990	I	pe	1953	0.0199	0.0184	Kaluzny & Pojmanski (1984b)
2445294.4760	I	pg	2032	0.0240	-	Braune et al. (1983)
2445295.2840	I	pe	2034	0.0200	0.0181	Kurpinska-Winiarska & Zakrzewski (1990)

Table 2—Continued

Hel.JD	Type	Method	E	$(O - C)$ Eq.2	$(O - C)$ Eq.7	Reference
2445296.2990	II	vis	2036.5	0.0200	-	Locher (1983a)
2445298.5360	I	pg	2042	0.0241	-	Braune et al. (1983)
2445323.2960	I	pe	2103	0.0189	0.0166	Pohl et al. (1983)
2445335.2840	II	vis	2132.5	0.0303	-	Locher (1983a)
2445345.2310	I	vis	2157	0.0307	-	Locher (1983a)
2445346.2380	II	vis	2159.5	0.0227	-	Locher (1983a)
2445359.2270	II	vis	2191.5	0.0202	-	Locher (1983a)
2445388.2590	I	vis	2263	0.0241	-	Locher (1983b)
2445576.4450	II	vis	2726.5	0.0355	-	Isles (1985a)
2445577.4580	I	vis	2729	0.0335	-	Isles (1985a)
2445587.3860	II	vis	2753.5	0.0149	-	Isles (1985a)
2445593.4817	II	pe	2768.5	0.0208	0.0149	Kurpinska-Winiarska & Zakrzewski (1990)
2445605.8630	I	pe	2799	0.0195	0.0134	Faulkner (1986)
2445609.5174	I	pe	2808	0.0200	0.0139	Kurpinska-Winiarska & Zakrzewski (1990)
2445621.2890	I	vis	2837	0.0180	-	Locher (1983c)
2445623.3330	I	vis	2842	0.0321	-	Hubscher & Mundry 1984
2445635.3060	II	vis	2871.5	0.0285	-	Isles (1985a)
2445635.2940	II	vis	2871.5	0.0165	-	Locher (1983c)
2445641.3810	II	vis	2886.5	0.0137	-	Locher (1983c)
2445651.3318	I	pe	2911	0.0178	0.0112	Pohl et al. (1985)
2445674.2680	II	vis	2967.5	0.0158	-	Locher (1984a)
2445681.3790	I	vis	2985	0.0220	-	Isles (1985a)
2445698.2264	II	pe	3026.5	0.0210	0.0137	Kurpinska-Winiarska & Zakrzewski (1990)
2445701.2650	I	vis	3034	0.0147	-	Locher (1984a)
2445731.3140	I	vis	3108	0.0207	-	Isles (1985b)
2445772.3060	I	vis	3209	0.0081	-	Locher (1984b)
2445943.8485	II	pe	3631.5	0.0214	0.0109	Faulkner (1986)
2445988.2970	I	vis	3741	0.0144	-	Locher (1984c)
2446001.3070	I	vis	3773	0.0328	-	Isles (1985b)
2446005.3520	I	vis	3783	0.0180	-	Locher (1984c)
2446021.5914	I	pe	3823	0.0179	0.0064	Faulkner (1986)
2446059.3650	I	vis	3916	0.0348	-	Isles (1985b)
2446113.3600	I	vis	4049	0.0336	-	Isles (1986)
2446114.3580	II	vis	4051.5	0.0167	-	Isles (1986)
2446321.6210	I	vis	4562	0.0237	-	Locher (1985)
2446327.4994	II	pe	4576.5	0.0152	-0.0003	Pohl et al. (1987)
2446351.4780	II	vis	4635.5	0.0407	-	Isles (1986)

Table 2—Continued

Hel.JD	Type	Method	E	$(O - C)$ Eq.2	$(O - C)$ Eq.7	Reference
2446355.3280	I	vis	4645	0.0338	-	Locher (1985)
2446355.3070	I	vis	4645	0.0128	-	Hubscher et al. (1986)
2446383.3280	I	vis	4714	0.0207	-	Isles (1986)
2446403.4110	II	vis	4763.5	0.0074	-	Locher (1986)
2446421.2970	II	vis	4807.5	0.0300	-	Locher (1986)
2446422.2880	I	vis	4810	0.0061	-	Hubscher et al. (1986)
2446440.5677	I	pe	4855	0.0164	-0.0007	Faulkner (1986)
2446688.4660	II	vis	5465.5	0.0600	-	Isles (1988)
2446714.6062	I	pe	5530	0.0141	-0.0066	Kurpinska-Winiarska & Zakrzewski (1990)
2446716.4360	II	pe	5534.5	0.0170	-0.0037	Kurpinska-Winiarska & Zakrzewski (1990)
2446745.4636	I	pe	5606	0.0166	-0.0046	Kurpinska-Winiarska & Zakrzewski (1990)
2446760.2930	II	vis	5642.5	0.0275	-	Locher (1987a)
2446761.2955	I	pe	5645	0.0150	-0.0063	Kurpinska-Winiarska & Zakrzewski (1990)
2446843.3230	I	vis	5847	0.0333	-	Locher (1987b)
2447068.4410	II	vis	6401.5	0.0319	-	Hubscher & Lichtenknecker (1988)
2447077.3660	II	vis	6423.5	0.0252	-	Hubscher & Lichtenknecker (1988)
2447088.3410	II	vis	6450.5	0.0385	-	Locher (1988a)
2447111.2870	I	vis	6507	0.0463	-	Locher (1988b)
2447113.3050	I	vis	6512	0.0344	-	Locher (1988c)
2447118.3640	II	vis	6524.5	0.0186	-	Locher (1988a)
2447118.7666	II	pe	6525.5	0.0152	-0.0109	Rainger et al. (1992)
2447119.7814	I	pe	6528	0.0150	-0.0111	Rainger et al. (1992)
2447128.3450	I	vis	6549	0.0529	-	Locher (1988c)
2447141.3010	I	vis	6581	0.0173	-	Locher (1988a)
2447143.3294	I	pe	6586	0.0158	-0.0106	Kurpinska-Winiarska & Zakrzewski (1990)
2447145.3640	I	vis	6591	0.0205	-	Locher (1988a)
2447153.2780	II	vis	6610.5	0.0177	-	Locher (1988a)
2447157.3540	II	vis	6620.5	0.0339	-	Locher (1987b)
2447206.2530	I	pe	6741	0.0115	-0.0157	Keskin & Pahi (1989)
2447207.2750	II	vis	6743.5	0.0186	-	Hubscher & Lichtenknecker (1988)
2447208.2920	I	vis	6746	0.0206	-	Locher (1988c)
2447439.4958	II	pe	7315.5	0.0152	-0.0151	Kurpinska-Winiarska & Zakrzewski (1990)
2447444.5695	I	pe	7328	0.0141	-0.0163	Kurpinska-Winiarska & Zakrzewski (1990)
2447449.4417	I	pe	7340	0.0144	-0.0160	Kurpinska-Winiarska & Zakrzewski (1990)
2447450.4569	II	pe	7342.5	0.0147	-0.0158	Kurpinska-Winiarska & Zakrzewski (1990)
2447452.4872	II	pe	7347.5	0.0150	-0.0154	Kurpinska-Winiarska & Zakrzewski (1990)
2447511.3531	II	pe	7492.5	0.0129	-0.0183	Keskin & Pahi (1989)

Table 2—Continued

Hel.JD	Type	Method	E	($O - C$) Eq.2	($O - C$) Eq.7	Reference
2447512.3675	I	pe	7495	0.0124	-0.0189	Hanzl (1990)
2447523.3790	I	vis	7522	0.0622	-	Locher (1989b)
2447523.3120	I	vis	7522	-0.0048	-	Locher (1989a)
2447524.3480	II	vis	7524.5	0.0163	-	Locher (1989a)
2447525.3420	I	vis	7527	-0.0047	-	Locher (1989a)
2447534.3010	I	vis	7549	0.0226	-	Locher (1989b)
2447561.2904	II	pe	7615.5	0.0139	-0.0180	Kurpinska-Winiarska & Zakrzewski (1990)
2447565.3460	II	vis	7625.5	0.0097	-	Locher (1989b)
2447823.5577	II	pe	8261.5	0.0141	-0.0213	Kurpinska-Winiarska & Zakrzewski (1990)
2447828.4289	II	pe	8273.5	0.0134	-0.0220	Kurpinska-Winiarska & Zakrzewski (1990)
2447834.3148	I	pe	8288	0.0125	-0.0230	Kurpinska-Winiarska & Zakrzewski (1990)
2447848.3226	II	pe	8322.5	0.0138	-0.0219	Kurpinska-Winiarska & Zakrzewski (1990)
2448501.5500	II	pe	9931.5	0.0093	-0.0351	Hubscher, Agerer & Wunder (1992)
2448569.1482	I	pe	10098	0.0107	-0.0345	Liu et al. (1993)
2448570.1604	II	pe	10100.5	0.0080	-0.0373	Liu et al. (1993)
2448604.2656	II	pe	10184.5	0.0103	-0.0354	Hubscher, Agerer & Wunder (1992)
2448623.3495	II	pe	10231.5	0.0129	-0.0331	Diethelm (1992)
2448928.4421	I	pe	10983	0.0068	-0.0432	Muyesseoglu, Gurol & Selam (1996)
2448987.3137	I	pe	11128	0.0104	-0.0404	Hubscher, Agerer & Wunder (1993)
2449287.5387	II	pe	11867.5	0.0085	-0.0462	Ogooza (1995)
2449341.3288	I	pe	12000	0.0054	-0.0500	Muyesseoglu, Gurol & Selam (1996)
2449342.3464	II	pe	12002.5	0.0081	-0.0474	Muyesseoglu, Gurol & Selam (1996)
2449625.3171	II	pe	12699.5	0.0063	-0.0529	Muyesseoglu, Gurol & Selam (1996)
2449625.5210	I	pe	12700	0.0072	-0.0520	Muyesseoglu, Gurol & Selam (1996)
2449688.4452	I	pe	12855	0.0035	-0.0565	Ogooza (1995)
2449689.4625	II	pe	12857.5	0.0059	-0.0542	Ogooza (1995)
2450043.2780	I	pe	13729	0.0043	-0.0604	Hegedus et al. (1996)
2450043.4802	II	pe	13729.5	0.0035	-0.0612	Hegedus et al. (1996)
2450045.5113	II	pe	13734.5	0.0047	-0.0601	Hegedus et al. (1996)
2450393.4395	II	pe	14591.5	0.0027	-0.0667	Selam, Gurol & Muyesseoglu (1999)
2450398.1064	I	CCD	14603	0.0007	-0.0687	Kim et al. (1997)
2450412.1138	II	CCD	14637.5	0.0016	-0.0680	Kim et al. (1997)
2450414.9553	II	CCD	14644.5	0.0012	-0.0685	Kim et al. (1997)
2450415.1583	I	CCD	14645	0.0012	-0.0685	Kim et al. (1997)
2450429.1652	II	CCD	14679.5	0.0016	-0.0683	Kim et al. (1997)
2450430.9914	I	CCD	14684	0.0008	-0.0690	Kim et al. (1997)
2450432.0068	II	CCD	14686.5	0.0013	-0.0686	Kim et al. (1997)

Table 2—Continued

Hel.JD	Type	Method	E	($O - C$) Eq.2	($O - C$) Eq.7	Reference
2450433.0213	I	CCD	14689	0.0008	-0.0691	Kim et al. (1997)
2450671.5357	II	pe	15276.5	-0.0018	-0.0748	Ogooza (1997)
2450725.3310	I	CCD	15409	0.0004	-0.0734	Diethelm & Blattler (1998)
2450729.3866	I	pe	15419	-0.0039	-0.0777	Selam, Gurol & Muyesseoglu (1999)
2450763.2888	II	pe	15502.5	-0.0016	-0.0758	Diethelm & Blattler (1998)
2451194.2393	I	CCD	16564	-0.0055	-0.0855	Blattler (1999)
2451470.3092	I	CCD	17244	-0.0063	-0.0899	Kim et al. (2003)
2451515.1711	II	CCD	17354.5	-0.0059	-0.0901	Kim et al. (2003)
2451542.9807	I	CCD	17423	-0.0063	-0.0909	Kim et al. (2003)
2451579.7217	II	CCD	17513.5	-0.0071	-0.0921	Nelson (2001)
2451928.2587	I	pe	18372	-0.0093	-0.0990	Pribulla et al. (2001)
2452528.5036	II	CCD	19850.5	-0.0152	-0.1128	Pribulla et al. (2002)
2452547.3805	I	CCD	19897	-0.0166	-0.1145	Borkovits et al. (2003)
2452322.2640	II	CCD	19342.5	-0.0137	-0.1086	Agerer & Hubscher (2003)
2452535.4050	II	CCD	19867.5	-0.0155	-0.1132	Agerer & Hubscher (2003)
2452576.4098	II	CCD	19968.5	-0.0153	-0.1136	Agerer & Hubscher (2003)
2452617.4140	II	CCD	20069.5	-0.0158	-0.1145	Agerer & Hubscher (2003)
2452677.2976	I	CCD	20217	-0.0151	-0.1147	Agerer & Hubscher (2003)
2452888.4068	I	CCD	20737	-0.0188	-0.1212	Krajci (2005)
2452908.5026	II	CCD	20786.5	-0.0193	-0.1219	Hubscher (2005)
2452992.3382	I	CCD	20993	-0.0199	-0.1236	Hubscher (2005)
2453003.2999	I	CCD	21020	-0.0198	-0.1237	Hubscher (2005)
2453028.2681	II	CCD	21081.5	-0.0198	-0.1240	Hubscher et al. (2005)
2453330.3190	II	CCD	21825.5	-0.0227	-0.1309	Hubscher et al. (2006)
2453409.2829	I	CCD	22020	-0.0231	-0.1324	Hubscher et al. (2006)
2453681.2900	I	CCD	22690	-0.0268	-0.1397	Hubscher et al. (2006)
2453683.3202	I	CCD	22695	-0.0266	-0.1394	Hubscher et al. (2006)
2453707.2731	I	CCD	22754	-0.0268	-0.1400	Hubscher et al. (2006)
2453764.3131	II	CCD	22894.5	-0.0279	-0.1419	Hubscher et al. (2006)
2454056.4176	I	CCD	23614	-0.0306	-0.1484	Biro et al. (2007)
2454033.8860	II	CCD	23558.5	-0.0299	-0.1474	Nelson (2007)
2453763.2983	I	CCD	22892	-0.0278	-0.1417	Hubscher & Walter (2007)
2454116.2986	I	CCD	23761.5	-0.0325	-0.1511	Hubscher & Walter (2007)
2454448.1874	I	CCD	24579	-0.0375	-0.1605	the paper
2454449.2042	II	CCD	24581.5	-0.0357	-0.1587	the paper

Table 3: Photometric solutions for SS Ari.

Parameters	Photometric elements	errors
	spotted solutions	
$g_1 = g_2$	0.32	assumed
$A_1 = A_2$	0.50	assumed
$x_{1R} = x_{2R}$	0.491	assumed
T_1	5860K	assumed
q	3.250	assumed
Ω_{in}	6.9452	–
Ω_{out}	6.3227	–
T_2	5488K	± 12 K
i	77.306	± 0.205
$L_1/(L_1 + L_2)(R)$	0.2814	± 0.0114
$\Omega_1 = \Omega_2$	6.8869	± 0.0093
$r_1(pole)$	0.2669	± 0.0006
$r_1(side)$	0.2785	± 0.0007
$r_1(back)$	0.3142	± 0.0013
$r_2(pole)$	0.4576	± 0.0006
$r_2(side)$	0.4925	± 0.0008
$r_2(back)$	0.5190	± 0.0071
f	9.4 %	± 0.8 %
θ ($^\circ$)	29.21	
ψ ($^\circ$)	284.50	
Ω ($^\circ$)	12.755	
T_s/T_*	0.8333	
$\sum \omega_i(O - C)_i^2$	0.0029038	

Table 4: The comparison of photometric solutions for SS Ari.

Parameters	Photometric elements	Photometric elements	Photometric elements
	unspotted solutions	unspotted solutions	spotted solutions
	Lu 1991	Liu et al. 1993	Kim et al. 2003
$g_1 = g_2$	0.32	0.32	0.32
$A_1 = A_2$	0.50	0.50	0.50
$x_{1B} = x_{2B}$	0.74	0.700	–
$x_{1V} = x_{2V}$	0.61	0.560	–
T_1	5950K	6050K	6144K
q	3.308	3.030	3.255
T_2	5745K	5908K	5860K
i	75.34	74.73	74.62
$L_1/(L_1 + L_2)(B)$	0.2902	0.2958	–
$L_1/(L_1 + L_2)(V)$	0.2835	0.2911	–
$\Omega_1 = \Omega_2$	6.9384	6.551	6.838
$r_1(pole)$	0.2672	0.2755	0.2706
$r_1(side)$	0.2789	0.2880	0.2828
$r_1(back)$	0.3160	0.3266	0.3218
$r_2(pole)$	0.4604	0.4553	0.4612
$r_2(side)$	0.4962	0.4900	0.4974
$r_2(back)$	0.5231	0.5182	0.5252
f	13.2 %	10.6 %	18.3 %

Table 5: The masses and orbital radii of the assumed third body in SS Ari. The calculated value about M_3 and a_3 are effective numbers.

Parameters	Third body			Units
	eccentric adding cycle		quadratic adding cycle	
A_3	0.1045	0.1045	0.0112	day
T_3	87.0	87.0	37.75	yr
e'	0.3948	0.3948	0	
w'	$-26^\circ.983$	$-26^\circ.983$	0°	
$a_{12}\sin i'$	14.5566	33.5485	1.9392	AU
$f(m)$	0.4075	4.9886	5.1172×10^{-3}	$M_\odot$
$m_3(i' = 90^\circ)$	1.66	7.50	0.271	$M_\odot$
$m_3(i' = 77.306^\circ)$	1.73	8.62	0.278	$M_\odot$
$a_3(i' = 90^\circ)$	26.2	15.9	10.1	AU
$a_3(i' = 77.306^\circ)$	26.5	16.5	10.1	AU




Fig. 1.— CCD photometric light curves in R band of SS Ari obtained by 60cm reflecting telescope on December 13, 14, 2007.




Fig. 2.— $(O - C)$ diagram of SS Ari formed by all available observations. The $(O - C)$ values were computed by using linear ephemeris (Eq.2). Solid cycles are referred to the photoelectric and CCD primary minima while open ones to the photoelectric and CCD secondary minima. Solid triangles denote visual and photographic primary minima, open triangles denote visual and photographic secondary minima, these data may not be precise. Solid line represents an eccentric ephemeris variation (Eq.3).




Fig. 3.— The residuals for the eccentric ephemeris (Eq.3). The symbols are the same as Figure 2.




Fig. 4.— Diagram of SS Ari formed by all available photoelectric and CCD observations. The $(O - C)$ values were computed by using a new determined linear ephemeris (Eq.7). The symbols are the same as Figure 2. Solid line represents a quadratic superimposed a cycle ephemeris variations (Eq.9), meanwhile dash line represent the quadratic variations (Eq.8).




Fig. 5.— $(O - C)_2$ values for SS Ari with respect to the cyclic ephemeris in Eq.(9). The symbols are the same as figure 2. Solid line refers to the theoretical orbit of an assumed third body.




Fig. 6.— The residuals for the quadratic superimposed a cycle ephemeris and a cyclic variations (Eq.9). The symbols are the same as Figure 2.




Fig. 7.— The relation between q and Σ for SS Ari. It hard to find a accurate mass ratio from the picture.




Fig. 8.— Observed (open circles) and theoretical (solid lines) light curves in R band for SS Ari, with a dark spot in the more massive component.




Fig. 9.— Geometrical structure of the shallow contact binary SS Ari with a dark spot on the more massive component at phase 0.00, 0.25, 0.50 and 0.75.